

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

6-4-2007

Soybean aphids found in Minnesota and Wisconsin

Marlin E. Rice

Iowa State University, merice@iastate.edu

Matthew E. O'Neal

Iowa State University, oneal@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

Rice, Marlin E. and O'Neal, Matthew E., "Soybean aphids found in Minnesota and Wisconsin" (2007). *Integrated Crop Management News*. Paper 1048.

<http://lib.dr.iastate.edu/cropnews/1048>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact digirep@iastate.edu.

INTEGRATED CROP MANAGEMENT

Search

Get the latest research-based information on crops. [Sign up to be notified](#) when new content is available!

ICM > 2007 > IC-498(13) -- June 4, 2007

Current Newsletter

You are viewing **archives** for the newsletter from 1993-2007. For current news, see [Integrated Crop Management News](#).

Archives 1993-2007



Announcements



Crop Production



Insects and Mites



Pesticide Education



Plant Diseases



Soils



Weed Management

[Image Gallery](#)

Printable Version

Printable version of this page

Related Articles

Aphids attack!
August 6, 2007

Extension agronomists update soybean aphid situation
July 30, 2007

Newly revised aphids publication available
July 23, 2007

Mid-July soybean aphid update

Soybean aphids found in Minnesota and Wisconsin

by Marlin E. Rice and Matt O'Neal, Department of Entomology

Our entomological colleagues to the north and northeast are reporting the first soybean aphids of the summer on seedling soybean plants.



Winged soybean aphids fly from buckthorn to soybean where they give birth to live aphids.
(Marlin E. Rice)

Eileen Cullen, extension entomologist, University of Wisconsin, reported that in a research plot near Madison on May 24, 8 of 60 plants were infested with 1-13 aphids per plant. Similar populations were found near Arlington on May 30. David Ragsdale, research entomologist, University of Minnesota, reported soybean aphids on V1-stage soybeans in east-central Minnesota during the week of May 28. Some fields had more than 25 percent of the plants infested with an average density of 5 aphids per plant. Furthermore, Chris DiFonzo, extension entomologist, Michigan State University, reported three research farms with infestations of soybean aphids. In Michigan, all sites had less than 1 percent of plants infested with aphids.

In Iowa in recent years, soybean aphids have been found during the last week of May in central or northeastern Iowa, but as of June 4, we have not yet received any reports of aphids in Iowa soybeans. Our aphid suction trap network with traps located near Ames, Chariton, Nashua, and Sutherland did not collect any winged soybean aphids the weeks preceding May

July 16, 2007

Soybean aphids exceed the economic threshold in northeast Iowa

July 9, 2007

Monitor soybean aphid populations on PIPE

July 2, 2007

Soybean aphid numbers increase...and decrease

July 2, 2007

Early soybean aphid outbreak--to the east

June 11, 2007

PIPE: Pest Information Platform for Extension and Education

March 26, 2007

Purdue University identifies the #1 predator of soybean aphids

March 26, 2007

18 and May 25. This may suggest that the Iowa population is small this year or maybe not because the traps won't necessarily represent the entire state. Also, it may take us several years to be able to interpret what the spring soybean aphid trap captures mean in relationship to early summer infestations in soybean fields.

The important point to remember is that soybean aphids will appear somewhere in Iowa in damaging numbers this summer--and field scouting will be necessary to determine the need for chemical control.

As of yet, no damaging populations occur in Iowa, but field scouting should start in 2-3 weeks so that informed decisions can be made in a timely manner.

Marlin E. Rice is a professor of entomology with extension and research responsibilities. Matt O'Neal is an assistant professor of entomology with research and extension responsibilities in field crops.

This article originally appeared on page 167 of the IC-498(13) -- June 4, 2007 issue.

Updated 06/06/2007 - 11:09am